

# Atlantic Multidecadal Variability and hydrological cycle in the Caspian Sea watershed

V.A. Semenov<sup>1,2</sup>, N.G. Nikitina<sup>1,2</sup>, I.I. Mokhov<sup>1</sup>

<sup>1</sup>A.M. Obukhov Institute of Atmospheric Physics RAS, Moscow, Russia

<sup>2</sup>Lomonosov Moscow State University, Moscow, Russia

[vasemenov@mail.ru](mailto:vasemenov@mail.ru)

The Caspian Sea level (CSL) experiences large variations that highly impact society and environment. The CSL variations are determined by water cycle changes integrated over the huge (more than 3 million square km) watershed area. Volga River discharge (VRD) explains more than 80% of the CSL variations during the last century. Many different factors have been previously proposed to contribute to VRD change including major models of natural climate variability, e.g., NAO [Rodionov 1994] and ENSO [Arpe et al. 2000], and anthropogenic forcing [e.g., Arpe et al. 1999]. Recently, it has been suggested that Atlantic Multidecadal Variability (AMV), the major low-frequency climate variability mode in the North Atlantic, may strongly affect even the global climate variations [e.g., Semenov et al. 2010]. Analysis of coupled climate model simulations revealed a significant link between AMV and hydrological cycle over Eurasia on inter-decadal time scale [Mokhov et al. 2008]. Here, we focus on the AMV impact on hydrological cycle in the Caspian Sea watershed using global model simulations.

The simulations are performed using ECHAM5 atmosphere general circulation model of T31 spectral resolution (appr. 3.8° lat/lon) coupled to thermodynamical mixed layer (50 m) ocean model. The coupled model is forced by periodically (60-yr) time-varying anomalous oceanic heat convergence flux in the North Atlantic and Arctic representing AMV. The anomalous heat flux pattern, model and experimental design are described in [Semenov et al. 2010].




Figure 1: Correlations between simulated annual AMV SST index (see the text) and precipitation-minus-evaporation for realistic (a) and doubled (b) forcing experiments. Data have been smoothed with 15-yr running averages.

Two simulations of 500 yrs duration are performed. One uses realistic AMV amplitude (about 0.09 PW integrated heat transport). Another uses a doubled amplitude of the anomalous heat flux pattern. These simulations are idealized as they do not involve ocean dynamics and other internal variability modes essentially interacting with the AMO. From the other hand, such experiments allow one a clear separation of the AMV-related signal that is the only time-varying external forcing factor in the coupled system. The experiments start

from the present-day climate conditions (2001-2010) and maximum of the AMV forcing.




Figure 2: Simulated annual Volga River discharge anomalies (VRD, in km<sup>3</sup>/yr, red), AMV SST index (in K, black) and applied integrated heat flux (HF) forcing (in PW, green) for realistic (a) and doubled (b) forcing experiments. All time series are 15-yr running averages.

The results reveal a significant positive correlation between AMV, as represented by averaged sea surface temperature anomalies in [50W-10W,40N-60N] region, and precipitation-minus-evaporation difference over the Eastern Eurasia including Caspian Sea watershed (Fig. 1). Simulation with realistic forcing exhibits weaker correlations. However, area of stronger correlation is found in the eastern part of the Volga River watershed. Correlation of AMV SST index with precipitation is stronger for both simulations. Simulated VRD anomalies are shown in Fig. 2 together with AMV SST index and applied heat flux forcing. In doubled forcing simulation, variations of VRD are closely linked with the AMV SST index (Fig. 2b). The link is not robust in the realistic forcing simulation where periods of strong correlation (e.g. 2000-2250) and weak correlation (e.g., 2250-2350) alternate (Fig. 2a).

An estimate of the interdecadal variations of the annual Volga river discharge related to the AMV (estimated from linear regression) amounts to about 40 cubic kilometers per year which is comparable to the observed changes during the 20<sup>th</sup> century. This indicates a possible role of the AMV in driving CSL changes. The results imply a potential decadal predictability for CSL.

This work was supported by the Russian Foundation for Basic Research; the Presidium of the Russian Academy of Sciences (programs 31, 4); the Ministry for Education and Science of the Russian Federation (state contracts nos. 11.519.11.5006 and 11.G34.31.0007); and the Russian Academy of Sciences (contract no. 74\_OK/11\_4).

## References

- Arpe, K., Bengtsson, L., Golitsyn, G. S., Mokhov, I. I., Semenov, V. A., and P. V. Sporyshev (1999): Doklady Earth Sciences, 366, No. 4, 552–556.
- Arpe, K., Bengtsson, L., Golitsyn, G. S., Mokhov, I. I., Semenov, V. A., and P. V. Sporyshev (2000): Connection between Caspian Sea level variability and ENSO. Geophys. Res. Lett., 27, 2693-2696.
- Mokhov, I.I., Semenov, V.A., Khon, V.C., Latif, M., and E. Roeckner (2008): Connection between Eurasian and North Atlantic Climate anomalies and natural variations in the Atlantic Thermohaline Circulation based on long-term model calculations. Doklady Earth Sciences, 419A, No. 3, 502–505
- Rodionov, S.N. Global and regional climate interaction: the Caspian Sea experience. 241 pp. Kluwer Academic Publ., Dordrecht, The Netherlands, 1994.
- Semenov, V.A., Latif, M., Dommenges, D., Keenlyside, N.S., Strehz, A., Martin, T., Park, W. (2010): The Impact of North Atlantic-Arctic Multidecadal Variability on Northern Hemisphere Surface Air Temperature. J. Climate 23, 5668-5677.